

# Herbicide

## CAUTION

# Weedeath

Cyndan\*

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**  
Active Constituents: 250g/L Amitrole, 220g/L Ammonium Thiocyanate

GROUP **F** HERBICIDE

FOR THE CONTROL OF WEEDS AND GRASSES IN GENERAL AND INDUSTRIAL SITUATIONS AS STATED IN THE DIRECTIONS FOR USE TABLE

SITUATION	WEED CONTROLLED	STATE	RATE Per 100L Knapsack	RATE Per 15L Knapsack	CRITICAL COMMENTS
Land - non - agricultural areas, around buildings, commercial/industrial, rights of way.	Achillea thistle, Common heliotrope, Couch, Creeping knapweed, Dandelion, Green polka-dot plant, Knapweed, Kikuyu grass, Onion weed, Sorrel, Sourgrass, Sticky cape gooseberry, Watsonia, Yarranilla top.	All States	1.1 Litres	185mL	Apply when weeds are actively growing, immediately prior to flowering. Respraying will be necessary to destroy regrowth and seedlings.
	African daisy, Apple-of-Sodom, Feenel, Hoary cress, Lilybird thistle, Blackberry, Gorse, Ox-eyed daisy	Vic, SA, Tas	2 Litres	300mL	Apply when weeds are actively growing, immediately prior to flowering.
Drains and margins of waterways	Common reed, Cumbungi, Nutgrass	All States	2.3 Litres	345mL	Apply twice. First treatment Dec. or Jan. and regrowth sprayed in autumn. Respraying in the following year is essential. DO NOT apply to blackberry bushes carrying mature or developing fruit.
	Water couch		2.2 Litres OR 4.5 Litres	330mL OR 875mL	Apply during flowering period between Jan. and May. Retreatment of Nutgrass will be necessary.
	Water Hyacinth		280mL	18mL	Apply in Nov. to Dec. with a further application at the same rate 6-8 weeks later. Apply as a single treatment between Feb. and May. Apply immediately prior to flowering. If infestation is large, treat in sections to avoid saturation of water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

### GENERAL INSTRUCTIONS:

**MIXING AND APPLICATION:** Add the required quantity to half the required volume of clear water. Stir or agitate and then top up to spray volume. A suitable Cyndan surfactant may be added to the spray mixture at the recommended rates to improve the efficiency of this product. Thoroughly wet all surface by aiming to apply 1L/10 square metres (900-1100 litres per hectare) using a knapsack sprayer. Thoroughly clean spray equipment after use.

**Resistant Weeds Warning:** Cyndan WEEDEATH is a member of the Triazole group of herbicides. Cyndan WEEDEATH has the inhibitors of carotenoid biosynthesis mode of action. For weed resistance management, Cyndan WEEDEATH is a Group F herbicide. Some naturally-occurring weed biotypes resistant to Cyndan WEEDEATH and other inhibitors of carotenoid biosynthesis herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Cyndan WEEDEATH or other inhibitors of carotenoid biosynthesis herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Cydan Chemicals accepts no liability for any losses that may result from the failure of cyndan WEEDEATH to control resistant weeds.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:** This product is a non-selective herbicide and may damage crops, pastures, and some ornamentals. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands and pastures.

**PROTECTION OF LIVESTOCK:** Do not feed grass clippings from treated areas to poultry or livestock, low hazard to bees.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate dams, streams, rivers or waterways with the chemical or uses containers. DO NOT spray across open bodies of water. When treating aquatic plants, minimise entry of spray into water by treating only exposed parts of plants up to the point of runoff.

**STORAGE AND DISPOSAL:** Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before puncture and bury in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Do not inhale spray mist. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre, Phone 13 11 26 if swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using ipecac Syrup APF. Additional information is listed in the Material Safety Data Sheet.

NBA Approva No. 51994/0999

EXPIRY DATE:

(B)

CONTENTS:

LITRES